arXiv:1303.2749v1 [math.AG] 12 Mar 2013

A NEW INEQUALITY ON THE HODGE NUMBER $h^{1,1}$ OF ALGEBRAIC SURFACES

JUN LU, SHENG-LI TAN, FEI YU, AND KANG ZUO

Abstract. We get a new inequality on the Hodge number $h^{1,1}(S)$ of fibred algebraic complex surfaces S, which is a generalization of an inequality of Beauville. Our inequality implies the Arakelov type inequalities due to Arakelov, Faltings, Viehweg and Zuo, respectively.

1. Introduction and main results

For a compact complex Kähler surface S, we have several Hodge numbers

$$h^{0,1}(S) = h^{1,0}(S) = q(S), \quad h^{0,2}(S) = h^{2,0}(S) = p_g(S), \quad h^{1,1}(S).$$

However, the Hodge number $h^{1,1}$ is not well understood comparing with the others. Lefschetz's (1,1)-theorem tells us that the Néron-Severi group $\mathrm{NS}(S) = H^{1,1}(S) \cap H^2(S, \mathbb{Z})$. Denote by $\rho(S) = \operatorname{rank} \mathrm{NS}(S)$ the Picard number of S, i.e., the rank of $\mathrm{NS}(S)$. Then we have $h^{1,1}(S) \geq \rho(S)$.

Theorem 1.1 ([BPV04], Corollary 5.4)**.** *If a compact Kähler surface S does not admit any fibration with connected fibers over a curve of genus $b \geq 2$, then*

$$h^{1,1}(S) \geq 2q(S) - 1. \tag{1.1}$$

In what follows, we consider the case when S admits a fibration $f : S \to C$ over a smooth curve C of genus b. It is obvious that $q(S) \geq b$. We assume that the fibers are connected and the generic fiber is a smooth curve of genus g. Let $F_1, \cdots, F_s$ be all singular fibers of f and $\ell(F_i)$ be the number of irreducible components of F_i.

Theorem 1.2 ([Bea81], Lemma 2)**.**

$$h^{1,1}(S) \geq \rho(S) \geq 2 + \sum_{i=1}^{s} (\ell(F_i) - 1). \tag{1.2}$$

In this paper, we will generalize Beauville's inequality.

Theorem 1.3. *Let $f : S \to C$ be a genus g fiberation as above, $b = g(C)$. Then*

$$h^{1,1}(S) \geq 2(q(S) - b)b + 2 + \sum_{i=1}^{s} (\ell(F_i) - 1). \tag{1.3}$$

---

[1] This work was supported by the SFB/TR 45 Periods, Moduli Spaces and Arithmetic of Algebraic Varieties of the DFG (German Research Foundation).

[2] The first and the second named authors are also supported by NSFC, the Science Foundation of the EMC and the Foundation of Scientific Program of Shanghai.

[3] The first author is also supported by the Fundamental Research Funds for the Central Universities.

[4] *Key words and phrases.* Hodge number, Arakelov inequality, singular Jacobian, singular fiber.

As a consequence,

$$h^{1,1}(S) \geq 2bq(S) + 2 - 2b,$$

it is an analogue of (1.1) for fibered surfaces.

Let $\Gamma_1, \cdots, \Gamma_k$ be all irreducible components of a given fiber F, and $\widetilde{\Gamma}_i \to \Gamma_i$ be the normalization of Γ_i. $g(F) := \sum_{i=1}^{k} g(\widetilde{\Gamma}_i)$ is called the geometric genus of F. One can see that $g(F) \leq p_a(F) = g$.

Denote by $q_f := q(S) - b$ the relative irregularity of f. Beauville [Bea81] proves that for any fiber F,

$$g(F) \geq q_f \geq 0. \tag{1.4}$$

Let s_1 be the number of singular fibers satisfying $g(F) < g$, without loss of generality, we assume that $F_1, \cdots, F_{s_1}$ satisfy $g(F) < g$. In fact, s_1 is exactly the number of singular fibers of the associated family of Jacobians of the fibers. The new inequality on $h^{1,1}$ is related to the Arakelov inequality due to Viehweg and the last author.

Theorem 1.4. *With the notations as above, we have*

$$\begin{aligned} 2\deg f_*\omega_{S/C} = {} & (g - q_f)(2b - 2 + s_1) - \sum_{i=1}^{s_1}(g(F_i) - q_f) \\ & - \left(h^{1,1}(S) - 2q_f b - 2 - \sum_{i=1}^{s}(\ell(F_i) - 1)\right) + \sum_{i=1}^{s_1} N_{\overline{F}_i}. \end{aligned}$$

where $g \geq N_{\overline{F}_i} := g - p_a(\overline{F}_{i,\mathrm{red}}) \geq 0$, *and* $\overline{F}_i = \sigma^* F_i$ *is the normal crossing model of* F_i *obtained by a sequence of blowing-ups, see Sect. 2. In particular, if* F_i *is semistable, then* $N_{\overline{F}_i} = 0$.

As an application of the inequalities (1.3), (1.4) and Theorem 1.4, we have

Corollary 1.5. (Viehweg-Zuo [VZ06]) *Let* $f : S \to C$ *be a non-trivial semistable fibration. Let* s_1 *be the number of fibers with non-compact Jacobians. Then we have*

$$\deg f_*\omega_{S/C} \leq \frac{g - q_f}{2}(2b - 2 + s_1). \tag{1.5}$$

Furthermore, Viehweg and Zuo prove that the above inequality becomes equality if and only if the curve C is a Shimura curve in the moduli space of curves.

Since $s_1 \leq s$, the above result implies the classical Arakelov inequality as follows.

Corollary 1.6 (Arakelov [Ara71], Faltings [Fal83])**.**

$$\deg f_*\omega_{S/C} \leq \frac{g - q_f}{2}(2b - 2 + s).$$

In particular, we have a weaker inequality

$$\deg f_*\omega_{S/C} \leq \frac{g}{2}(2b - 2 + s). \tag{1.6}$$

Remark 1.7. In fact the inequality (1.6) is strict when $g \geq 2$. One can also get it by combining Cornalba-Harris-Xiao's inequality [CH88, Xia87]

$$\frac{4g - 4}{g}\deg f_*\omega_{S/C} \leq \omega_{S/C}^2$$

with Vojta's canonical class inequality [Voj88]

$$\omega_{S/C}^2 < (2g-2)\cdot(2b-2+s), \tag{1.7}$$

which is indeed strict (cf. [Tan95, Liu96]).

When $f : S \to \mathbb{P}^1$ is a semistable family over $\mathbb{P}^1$, the equality in Theorem 1.4 can be rewritten as

$$\begin{aligned}\frac{1}{2}(g-q(S))(s_1-4) = p_g(S) + \frac{1}{2}\sum_{i=1}^{s_1}(g(F_i)-q(S)) \\ + \frac{1}{2}\left(h^{1,1}(S) - 2 - \sum_{i=1}^{s}(\ell(F_i)-1)\right),\end{aligned} \tag{1.8}$$

where the formula $\deg f_*\omega_{S/C} = \chi(\mathcal{O}_S) - (g-1)(b-1)$ is used.

Corollary 1.8. *Assume that $f : S \to \mathbb{P}^1$ is semistable and non-trivial. Then we have $s_1 \geq 4$. $s_1 = 4$ if and only if the following conditions are satisfied.*

1. $p_g(S) = 0$*;* $g(F_i) = q(S)$ *for* $i = 1, \cdots, 4$*;*
2. $h^{1,1}(S) = 2 + \sum_{i=1}^{s}(\ell(F_i)-1)$.
3. $q(S) \leq 1$*;*

(1) and (2) are direct consequences of (1.2) and (1.4). (3) will be proved in §4. In §5, we will present two examples of genus $g = 2$ with $s_1 = 4$ and $q(S) = 1$.

In the case when f is non-semistable, we have

$$N_{\overline{F}_i} = g - p_a(\overline{F}_{i,\mathrm{red}}) \leq g - g(\overline{F}_i) = g - g(F_i) \leq g - q_f.$$

We get an inequality as follows.

Corollary 1.9. *If f is non-semistable, then*

$$\deg f_*\omega_{S/C} \leq (g-q_f)(b-1+s_1).$$

2. Proof of Theorem 1.3

Let $f : S \to C$ be a fibration of genus g over a smooth curve C of genus b, and $F_1, \cdots, F_s$ be all singular fibers. Due to Beauville's inequality (1.2), we can assume that $b > 0$ and $q_f > 0$. Consider $H^0(\Omega_S) = V_1 \oplus V_0$, where $V_0 = f^*H^0(\Omega_C)$ and $\dim V_1 = q_f$. Let

$$V_0 = \langle \alpha_1 \cdots \alpha_b \rangle, \quad V_1 = \langle \theta_1, \cdots, \theta_{q_f} \rangle,$$

where α_i's (resp. θ_j's) are the base of V_0 (resp. V_1) as a $\mathbb{C}$-vector space.

We define a homomorphism

$$h : V_0 \otimes \overline{V}_1 \oplus \overline{V}_0 \otimes V_1 \longrightarrow H^{1,1}(S),$$

by $h(x \otimes y) = x \wedge y$ for $x \otimes y \in V_0 \otimes \overline{V}_1 \oplus \overline{V}_0 \otimes V_1$.

Let V_2 be the subgroup of Pic(S)$\otimes\mathbb{R}$ generated by the classes of the components of all fibers. Chern class induces a homomorphism $c_1 : V_2 \to H^{1,1}(S) \cap H^2(S, \mathbb{R})$. By the semi-negativity of the intersection matrix of a fiber, we have

$$\dim \mathrm{Im}(c_1) = 1 + \sum_{i=1}^{s}(\ell(F_i)-1).$$

Lemma 2.1. *For any ample divisor H, $c_1(H) \notin \operatorname{Im} h + (\operatorname{Im} c_1) \otimes_{\mathbb{R}} \mathbb{C}$.*

Proof. Suppose that $c_1(H) = \alpha + \beta \in \operatorname{Im} h + (\operatorname{Im} c_1) \otimes_{\mathbb{R}} \mathbb{C}$ for some $\alpha \in \operatorname{Im} h$. Let F be a general fiber. For any $\alpha \in \operatorname{Im} h$, by the definition of h, one can see easily that $\alpha|_F = 0$. On the other hand, Zariski's lemma implies $\beta|_F = 0$ for any $\beta \in \operatorname{Im} c_1$. Hence $c_1(H)|_F = 0$, i.e., $HF = 0$, a contradiction. □

Note that $\alpha_i \wedge \bar{\alpha}_k$ is the pull-back of an element in $H^{1,1}(C)$ since $h^1(C, \Omega_C) = h^0(C, \mathcal{O}_C) = 1$. So we can assume that $\alpha_i \wedge \bar{\alpha}_k = \varepsilon_{ik} \alpha_1 \wedge \bar{\alpha}_1$ as cohomology classes in $H^{1,1}(C)$, where ε_{ik} are complex numbers.

Lemma 2.2. *The matrix $E = (\varepsilon_{ik})_{1 \le j, k \le b}$ is invertible.*

Proof. Suppose that there is a vector $(\lambda_1, \cdots, \lambda_b) \neq 0$, such that $E \cdot (\lambda_1, \cdots, \lambda_b)^T = 0$, i.e., $\sum\limits_k \varepsilon_{ik} \lambda_k = 0$ for all i. Then $\sum\limits_k \varepsilon_{ik} \lambda_k \alpha_1 \wedge \bar{\alpha}_1 = 0$. Namely, one has

$$\sum_k \lambda_k \alpha_i \wedge \bar{\alpha}_k = \alpha_i \wedge \sum_k \lambda_k \bar{\alpha}_k = 0, \quad \text{for all } i.$$

Therefore we get $\sum\limits_k \bar{\lambda}_k \alpha_k \wedge \sum\limits_k \lambda_k \bar{\alpha}_k = 0$, that is, $\sum\limits_k \bar{\lambda}_k \alpha_k = 0$. So $\lambda_i = 0$ for all i, a contradiction. □

Lemma 2.3. *h is injective.*

Proof. Suppose that there is a nonzero element in the kernel of h,

$$\sum_{j=1}^{q_f} \sum_{i=1}^{b} a_{ij} \alpha_i \wedge \bar{\theta}_j + \sum_{j=1}^{q_f} \sum_{i=1}^{b} b_{ij} \bar{\alpha}_i \wedge \theta_j = du. \tag{2.1}$$

(A Zero cohomology class means an exact form). Note that $d\alpha_k = d\theta_l = 0$, $\bar{\alpha}_i \wedge \bar{\alpha}_k = 0$. By wedging $\bar{\alpha}_k \wedge \theta_l$ on both sides, one gets

$$\sum_{j=1}^{q_f} \sum_{i=1}^{b} a_{ij} \alpha_i \wedge \bar{\theta}_j \wedge \bar{\alpha}_k \wedge \theta_l = d(u \wedge \bar{\alpha}_k \wedge \theta_l). \tag{2.2}$$

So

$$\sum_{j=1}^{q_f} \sum_{i=1}^{b} a_{ij} \varepsilon_{ik} \alpha_1 \wedge \bar{\alpha}_1 \wedge \bar{\theta}_j \wedge \theta_l = d(-u \wedge \bar{\alpha}_k \wedge \theta_l). \tag{2.3}$$

Let $\omega_k = \sum\limits_{j=1}^{q_f} \sum\limits_{i=1}^{b} \bar{a}_{ij} \bar{\varepsilon}_{ik} \theta_j$. We have $\alpha_1 \wedge \bar{\alpha}_1 \wedge \bar{\omega}_k \wedge \theta_l = d(-u \wedge \bar{\alpha}_k \wedge \theta_l)$. It implies that

$$\alpha_1 \wedge \bar{\alpha}_1 \wedge \bar{\omega}_k \wedge \omega_k = d(-u \wedge \alpha_k \wedge \omega_k).$$

By Stokes formula,

$$0 = \int_S \alpha_1 \wedge \bar{\alpha}_1 \wedge \bar{\omega}_k \wedge \omega_k = \int_S (\alpha_1 \wedge \omega_k) \wedge \overline{(\alpha_1 \wedge \omega_k)}\ .$$

So $\alpha_1 \wedge \omega_k = 0$, i.e., $\omega_k = f^* \beta_k$ for some $\beta_k \in H^0(C, \Omega_C)$. Thus $\omega_k \in V_0 \cap V_1$, i.e., for any k, $\omega_k = 0$. Hence

$$\sum_i a_{ij} \varepsilon_{ik} = 0, \qquad \text{for any } j \text{ and } k.$$

Therefore Lemma 2.2 implies that $a_{ij}=0$ for all i and j. Similarly, we have $b_{ij}=0$. It is a contradiction. □

Lemma 2.4. $\operatorname{Im} h \cap (\operatorname{Im} c_1)\otimes_{\mathbb{R}} \mathbb{C}=0$.

Proof. Note that $\operatorname{Im} c_1 \subseteq H^{1,1}(S)\cap H^2(S,\mathbb{R})$. Let $\Gamma_{i1}, \cdots, \Gamma_{i\ell_i}$ be the irreducible components of F_i, let $\omega = c_1(F)$ and let $\omega_{ij}=c_1(\Gamma_{ij})$. Assume that

$$x\cdot\omega+\sum_{i=1}^{s}\sum_{j=1}^{\ell_i-1}x_{ij}\cdot\omega_{ij}=t\in\operatorname{Im} h,\qquad x,\ x_{ij}\in\mathbb{C}. \tag{2.4}$$

Note that for any component Γ in the fibers, $t|_\Gamma=0$ because α_i's are pullback of forms on the base C. Similarly, $\omega|_\Gamma=0$.

$$\sum_{i=1}^{s}\sum_{j=1}^{\ell_i-1}x_{ij}\cdot\omega_{ij}|_\Gamma=0. \tag{2.5}$$

Thus

$$\sum_{i=1}^{s}\sum_{j=1}^{\ell_i-1}x_{ij}\cdot\int\omega_{ij}|_\Gamma=0. \tag{2.6}$$

Namely,

$$\sum_{i=1}^{s}\sum_{j=1}^{\ell_i-1}x_{ij}\cdot\Gamma_{ij}\Gamma=0\qquad\text{for any }\Gamma. \tag{2.7}$$

We know that the intersection matrix of Γ_{ij} is negative definite, for $1\le i\le s$ and $1\le j\le \ell_i-1$, we have $x_{ij}=0$. Thus we get that $x\cdot c_1(F)=t\in\operatorname{Im} h$ for some $x\in\mathbb{C}$.

If $x\neq 0$, then $c_1(F)\in\operatorname{Im} h$. Note that $c_1(F)=c\cdot\alpha_1\wedge\bar\alpha_1\neq 0$, i.e., $c\neq 0$. Let

$$c_1(F)=\sum_{j=1}^{q_f}\sum_{i=1}^{b}a_{ij}\alpha_i\wedge\bar\theta_j+\sum_{j=1}^{q_f}\sum_{i=1}^{b}b_{ij}\bar\alpha_i\wedge\theta_j\ ,$$

As forms, we have

$$\sum_{j=1}^{q_f}\sum_{i=1}^{b}a_{ij}\alpha_i\wedge\bar\theta_j+\sum_{j=1}^{q_f}\sum_{i=1}^{b}b_{ij}\bar\alpha_i\wedge\theta_j-c\cdot\alpha_1\wedge\bar\alpha_1=du. \tag{2.8}$$

Similar to the proof of the previous lemma, we get also (2.2). Use the same proof as above, we get $a_{ij}=b_{ij}=0$ for any i and j. Hence $-c\cdot\alpha_1\wedge\bar\alpha_1=du$. Thus as a class, $c_1(F)=c\cdot\alpha_1\wedge\bar\alpha_1=0$, a contradiction. This proves the lemma □

Combing the above lemmas, we have

$$h^{1,1}\ge 1+\dim\operatorname{Im} h+\dim\left((\operatorname{Im} c_1)\otimes_{\mathbb{R}}\mathbb{C}\right).$$

Then we get the desired inequality (1.3). We complete the proof of Theorem 1.3.

3. Proof of Theorem 1.4

Given a curve B in a surface X (non-zero effective divisor), we denote by $B_{\rm red}$ the reduced part of B. Let $\Gamma_1, \cdots, \Gamma_\ell$ be all irreducible components of B, and $\widetilde{\Gamma}_i \to \Gamma_i$ be the normalization. As in the introduction, we define

$$g(B) = \sum_{i=1}^{\ell} g(\widetilde{\Gamma}_i), \qquad N_B = p_a(B) - p_a(B_{\rm red}).$$

Let $q \in B$ be a singular point of $B_{\rm red}$, we denote by $\mu_q(B)$ the Milnor's number of $B_{\rm red}$ at q, and by $m_q(B)$ the multiplicity of $B_{\rm red}$ at q. Let $\mu_B = \sum\limits_{q\in B} \mu_q(B)$, where q runs over all singularities of $B_{\rm red}$.

Let $\sigma : \bar{S} \to S$ be the blowing-up at q, E the exceptional curve, and $\bar{B}$ the strict transform of B in $\bar{S}$. Assume that $\bar{B}$ intersects E at r points $q_1, \cdots, q_r$.

Lemma 3.1. *Assume that B is a reduced curve with a singular point $q \in B$. Then*

(1) $\chi_{\rm top}(B) = 2\chi(\mathcal{O}_B) + \mu_B$.

(2) *Let $m = m_q(B)$. Then*

$$\mu_q(B) = \sum_{i=1}^{r} \mu_{q_i}(\bar{B}) + m(m-1) - (r-1),$$

$$\mu_q(B) = \sum_{i=1}^{r} \mu_{q_i}(\bar{B}+E) + (m-1)(m-2) - 1,$$

(1) and (2) are proved in [Tan94] (Lemma 1.1 and Lemma 1.3.)

Definition 3.2. A *partial resolution* of the singularities of B is a sequence of blowing-ups $\sigma = \sigma_1 \circ \sigma_2 \circ \cdots \circ \sigma_r : \bar{X} \to X$

$$(\bar{X}, \sigma^* B) = (X_r, B_r) \xrightarrow{\sigma_r} X_{r-1} \xrightarrow{\sigma_{r-1}} \cdots \xrightarrow{\sigma_2} (X_1, B_1) \xrightarrow{\sigma_1} (X_0, B_0) = (X, B),$$

satisfying the following conditions:

(i) $B_{r,\rm red}$ has at worst ordinary double points as its singularities.

(ii) $B_i = \sigma_i^* B_{i-1}$ is the total transform of B_{i-1}.

Furthermore, σ is called the *minimal partial resolution* of the singularities of B if

(iii) σ_i is the blowing-up of X_{i-1} at a singular point $(B_{i-1,\rm red}, p_{i-1})$ which is not an ordinary double point for any $i \le r$. We denote by m_{i+1} the multiplicity of $(B_{\rm red,i}, p_i)$.

In what follows, we always assume that the partial resolutions are minimal, and we denote by $r = r(B)$ the number of blowing-ups in the minimal resolution and by $\overline{B} = B_r$. Note that

$$\alpha(B) = \mu_{\overline{B}} - \ell(\overline{B}) + 1$$

is the first Betti's number of the dual graph of $\overline{B}$ by Euler's formula. This number is determined uniquely by B.

From Lemma 3.1 and a straightforward computation, one gets the following lemma.

Lemma 3.3. (1) $\mu_{\overline{B}} = \mu_B - \sum\limits_{i=1}^{r} (m_i - 1)(m_i - 2) + r$.

(2) $N_{\overline{B}} = N_B + \frac{1}{2}\sum_{i=1}^{r}(m_i-1)(m_i-2)$.

(3) $p_a(\overline{B}_{\mathrm{red}}) = p_a(B_{\mathrm{red}}) - \frac{1}{2}\sum_{i=1}^{r}(m_i-1)(m_i-2)$.

For any fiber F, we define $e_F = \chi_{\mathrm{top}}(F_{\mathrm{red}}) - (2-2g)$.

Corollary 3.4. *(1) $e_F = 2N_F + \mu_F = 2N_{\overline{F}} + \mu_{\overline{F}} - r(F)$.*
(2) $p_a(\overline{F}_{\mathrm{red}}) \geq 0$ and $0 \leq N_F \leq N_{\overline{F}} \leq g$.
(3) $\alpha(F) = p_a(\overline{F}_{\mathrm{red}}) - g(F) \geq 0$ and $g = g(F) + N_{\overline{F}} + \alpha(F)$.

Proof. It follows from Lemma 3.1 (1) and Lemma 3.3 immediately. Note that $\overline{F}_{\mathrm{red}} = \Gamma_1 + \cdots + \Gamma_\ell$ is a connected nodal curve, $p_a(\Gamma_i) = g(\Gamma_i) + \alpha(\Gamma_i)$ and

$$p_a(\overline{F}_{\mathrm{red}}) = p_a(\Gamma_1) + \cdots + p_a(\Gamma_\ell) + \sum_{i<j}\Gamma_i\Gamma_j - \ell + 1 \geq 0.$$

$\alpha(F) = \alpha(\Gamma_1) + \cdots + \alpha(\Gamma_\ell) + \sum_{i<j}\Gamma_i\Gamma_j - \ell + 1 \geq 0$. □

Let $\omega_{S/C} := \omega_S \otimes (f_*\omega_C)^\vee$ be the relative canonical sheaf. The relative invariants of f are defined as follows.

$$\chi_f := \deg f_*\omega_{S/C}, \qquad K_f^2 := \omega_{S/C}\cdot\omega_{S/C}, \qquad e_f := \sum_F e_F.$$

It is well known that

$$\begin{cases} K_f^2 = c_1^2(S) - 8(g-1)(g(C)-1), \\ \chi_f = \chi(\mathcal{O}_S) - (g-1)(g(C)-1), \\ e_f = c_2(S) - 4(g-1)(g(C)-1). \end{cases}$$

By the definitions of e_f, χ_f and Hodge theory, one has

$$\begin{aligned} e_f &= 2 - 4q + 2p_g + h^{1,1}(S) - 4(g-1)(b-1), \\ \chi_f &= 1 - q + p_g - (g-1)(b-1). \end{aligned}$$

Thus $2\chi_f - e_f = 2q + 2(g-1)(b-1) - h^{1,1}(S)$, i.e.,

$$2\chi_f = e_f + (g-q_f)(2b-2) + 2q_f b + 2 - h^{1,1}(S). \tag{3.1}$$

On the other hand, from Corollary 3.4 and the fact that $r(F_i) = \ell(\overline{F}_i) - \ell(F_i)$, we have

$$\begin{aligned} e_f &= \sum_{i=1}^{s} e_{F_i} = \sum_{i=1}^{s}(2N_{\overline{F}_i} + \mu_{\overline{F}_i} - r(F_i)) = \sum_{i=1}^{s}(2N_{\overline{F}_i} + \mu_{\overline{F}_i} - \ell(\overline{F}_i) + \ell(F_i)) \\ &= \sum_{i=1}^{s} N_{\overline{F}_i} + \sum_{i=1}^{s}(g - p_a(\overline{F}_{i,\mathrm{red}}) + \alpha(F_i)) + \sum_{i=1}^{s}(\ell(F_i) - 1) \\ &= \sum_{i=1}^{s} N_{\overline{F}_i} + \sum_{i=1}^{s_1}(g - g(F_i)) + \sum_{i=1}^{s}(\ell(F_i) - 1) \\ &= \sum_{i=1}^{s} N_{\overline{F}_i} - \sum_{i=1}^{s_1}(g(F_i) - q_f) + \sum_{i=1}^{s}(\ell(F_i) - 1) + (g-q_f)s_1. \end{aligned} \tag{3.2}$$

Substitute (3.2) into (3.1), we obtain the equality in Theorem 1.4.

4. Applications

The following lemma is due to Beauville ([Bea81], Lemma 1). The original proof for the case when $C = \mathbb{P}^1$ works for the general case. For the reader's convenience, we would like to recall Beauville's proof.

Lemma 4.1. ([Bea81]) $g(F_i) \geq q_f$.

Proof. Let $\widetilde{F}$ be the normalization of F, and $\beta : J(\widetilde{F}) \to \mathrm{Alb}(S)$ be the natural map between the jacobian $J(\widetilde{F})$ and the Albanese variety $\mathrm{Alb}(S)$. Considering the abelian variety $Q = \mathrm{Alb}(S)/\mathrm{Im}\beta$, one gets an induced map $\bar{\alpha} : S \to Q$. $\bar{\alpha}(F)$ is a point in Q since $J(\widetilde{F}) \to Q$ is zero. Therefore, by the rigidity theorem, $\bar{\alpha}$ contracts all fibers of f. So $\bar{\alpha}$ has a factorization through f.

$$\begin{tikzcd}
S \arrow[r, "\bar{\alpha}"] \arrow[d, "f"'] & Q \\
C \arrow[ur] &
\end{tikzcd}$$

Since the image of S in $\mathrm{Alb}(S)$ generates $\mathrm{Alb}(S)$, we see that the image of C in Q generates Q. Thus we get a surjective map $v : J(C) \to Q$.

We have the following commutative diagram.

$$\begin{tikzcd}
\widetilde{F} \arrow[r] \arrow[d] & J(\widetilde{F}) \arrow[d, "\beta"'] \arrow[dr] & \\
S \arrow[r, "\alpha"] \arrow[d, "f"'] & \mathrm{Alb}(S) \arrow[r] \arrow[d, "g"] & Q \\
C \arrow[r, "j"'] & J(C) \arrow[ur, "v"'] &
\end{tikzcd}$$

If $g(C) = 0$, then Q is zero. Namely, $J(\widetilde{F}) \to \mathrm{Alb}(S)$ is surjective. Hence $g(F) = \dim J(\widetilde{F}) \geq \dim \mathrm{Alb}(S) = q$. If $g(C) > 0$, then $v : J(C) \to Q$ is surjective. Thus

$$g(C) = \dim J(C) \geq \dim Q = q - \dim \mathrm{Im}\,\beta,$$

i.e., $\dim \mathrm{Im}\,\beta \geq q_f$. So $g(F) = \dim J(\widetilde{F}) \geq \dim \mathrm{Im}\,\beta \geq q_f$. □

Similarly, from Corollary 3.4 and Lemma 4.1, we get Corollary 1.9.

Proof of Corollary 1.8:

From (1.8) and the inequalities, we only need to prove that $q \leq 1$.

Suppose that $q(S) \geq 2$. Because $p_g(S) = 0$, S is a ruled surface. The Albanese map $\alpha : S \to \mathrm{Alb}(S)$ induces the $\mathbb{P}^1$-fibration, and $B = \mathrm{Im}\,\alpha$ is a curve of genus q.

Because $g(F_i) = q \geq 2$, at least one irreducible component of $\overline{F}_i$, say Γ_1, doesn't lie in the fibers of α. So $g(\Gamma_1) \geq q = g(B)$ by Hurwitz formula. Thus $g(\Gamma_1) = q$ and Γ_1 is a section of $\alpha : S \to B$. It implies Γ_1 is the unique horizonal irreducible component of F_i since $g(F_i) = q$. Since F_i is semistable, $F_i = \Gamma_1 +$ component contracted by α. Let F' (resp. F) be a general fiber of α (resp. f). One has $F'F_i = F'\Gamma_1 = 1$, and hence $F'F = 1$. So $F \cong B$. Therefore f is isotrivial. Since

$f : S \to \mathbb{P}^1$ is semistable, f must be trivial, a contradiction. This completes the proof of Corollary 1.8. □

5. Examples

We will construct two semistable families $f : S \to \mathbb{P}^1$ with $s_1 = 4$, $g = 2$ and $q(S) = 1$.

Example 5.1 ([Xia85], Example 4.7)**.** Take six lines in $\mathbb{P}^2$ as follows.

$$P_1 : X = 0, \qquad P_2 : X - Y = 0, \qquad P_3 : Y = 0,$$
$$Q_1 : 2X + Y - Z = 0, \qquad Q_2 : X + Y - Z = 0, \qquad Q_3 : X + 2Y - Z = 0.$$



This configration of 6 lines has 4 triple points x, z_1, z_2, z_3, and 3 double points y_1, y_2 and y_3. Their coordinates are as follows. $x = [0, 0, 1]$ and

$$y_1 =[1/2, 0, 1], \qquad y_2 =[1/2, 1/2, 1], \qquad y_3 =[0, 1/2, 1],$$
$$z_1 =[1, 0, 1], \qquad z_2 =[1/3, 1/3, 1], \qquad z_3 =[0, 1, 1]$$

By Bertini's theorem, one can find an irreducible and reduced curve D of degree 4 in $\mathbb{P}^2$ satisfying the following conditions.

(1) D has ordinary double points at y_1, y_2 and y_3, and no other singular points.
(2) D passes through z_1, z_2, z_3 and x.

By blowing-up $\mathbb{P}^2$ at x, we get a ruled surface $\varphi : P \to \mathbb{P}^1$. Thus we can construct a double cover $\pi : X \to P$ branched along the curve $R = D + P_1 + P_2 + P_3 + Q_1 + Q_2 + Q_3$. The double cover gives us a semistable fibration $f : S \to \mathbb{P}^1$ of genus 2.

By a straightforward computation, we see that f is a Lefschetz pencil. Furthermore, we have $K_S^2 = -3$, $p_g(S) = 0$, $q(S) = 1$. Thus $K_f^2 = 5$, $\chi_f = 1$, $e_f = 7$. So f admits 7 singular fibers. $K_f^2 - 2\chi_f = 3$, this means that 3 singular fibers are not irreducible, so $s_1 = 7 - 3 = 4$.

$$s_1 = 4, \quad s = 7, \quad p_g = 0, \quad q = 1, \quad h^{1,1} = 5.$$

Example 5.2. Let $\mathrm{pr}_i : \mathbb{P}^1 \times \mathbb{P}^1 \to \mathbb{P}^1$ be the i-th projection and F_i^0 be a general fiber of pr_i ($i = 1, 2$). Let B_0 be a smooth irreducible curve of type $(2, 1)$, i.e., $B_0 \equiv 2F_1^0 + F_2^0$.



There are two fibers Γ_1^0 and Γ_2^0 of pr_2 tangent to B_0. By choosing two general fibers Γ_3^0 and Γ_4^0 of pr_2, one can construct a double cover $\pi_0 : E \times \mathbb{P}^1 \to \mathbb{P}^1 \times \mathbb{P}^1$ ramified over $B_{\pi_0} = \sum_{i=1}^{4} \Gamma_i^0$, where E is the pullback elliptic curve of a general fiber of pr_1.

Let

$$B_\pi := \pi_0^* B_0 \equiv 2F_1 + 2F_2, \qquad \pi_0^*\Gamma_i^0 = 2\Gamma_i, \qquad L = F_1 + \Gamma_3,$$

where F_i are general fibers of the i-th projection of $E \times \mathbb{P}^1$. Thus $B \equiv 2L$. We can construct a double cover $\pi : S_0 \to E \times \mathbb{P}^1$ branched along B_π. Thus we get a fibration $f : S \to \mathbb{P}^1$ of genus 2.

Now we claim that f has six singular fibers, and four of them have non-compact Jacobians. Let E_1 and E_2 be the elliptic fibers of $\mathrm{pr}_1 : E \times \mathbb{P}^1 \to \mathbb{P}^1$ such that the image fiber $\pi_0(E_i)$ passes through the tangent point $B_0 \cap \Gamma_i^0$ $(i = 1, 2)$. Let E_3, E_4 (resp. E_5, E_6) are other elliptic fibers whose image fibers pass through the intersection points of $B_0 \cap \Gamma_3^0$ (resp. $B_0 \cap \Gamma_4^0$). Take $p_i = E_i \cap \Gamma_i$, $p_{i+2} = E_{i+2} \cap \Gamma_3$, $p_{i+4} = E_{i+4} \cap \Gamma_4$ $(i = 1, 2)$ and $q_j = E_j \cap \Gamma_3$ $(j = 1, \cdots, 6)$.



Since $p_i \neq q_i$ and

$$2p_i = B_\pi\,|_{E_i} \equiv 2L|_{E_i} = 2q_i, \qquad i = 1, 2,$$

$\pi^{-1}(E_i)$ is irreducible $(i = 1, 2)$. So the fiber $\widetilde{F}_i$ of f corresponding to $\pi^{-1}(E_i)$ $(i = 1, 2)$ can be written as $\widetilde{F}_i = C_1 + C_2$ with $C_1 C_2 = 2$, where C_1 is a smooth elliptic curve and C_2 is a (-2)-curve.

$p_3 = q_3$ (resp. $p_4 = q_4$) implies that $\pi^{-1}E_3$ (resp. $\pi^{-1}E_4$) is reducible. So the corresponding fiber $\widetilde{F}_3$ (resp. $\widetilde{F}_4$) of f is a nodal curve $C_1 + C_2$ with $C_1 C_2 = 1$ where C_1, C_2 are smooth elliptic curves.

Similarly, one can check that $\pi^{-1}E_5$ and $\pi^{-1}E_6$ are irreducible. Thus the corresponding fiber $\widetilde{F}_5$ and $\widetilde{F}_6$ are singular elliptic curves with only one node.

By a straightforward computation, one has

$$s_1 = 4, \quad s = 6, \quad p_g = 0, \quad q = 1, \quad h^{1,1} = 6.$$

Remark 5.3. The third author proves in [Yu00] that if $s_1 = 4$ and $s = 5$, then $g = 2$.

Acknowledgements: The authors would like to thank Dr. Xin Lü for pointing out an error in the original proof of Lemma 2.4.

REFERENCES

[Ara71] S. Ju. Arakelov:*Families of algebraic curves with fixed degeneracies* (Russian), Izv. Akad. Nauk SSSR Ser. Mat. 35 (1971), 1269-1293.

[BPV04] W. Barth, K. Hulek, C. Peters, A. Van de Ven: *Compact Complex Surfaces, Second edition*, Springer-Verlag, Berlin, 2004.

[Bea81] A. Beauville: *Le nombre minimum de fibres singulières d'une courbe stable sur* $\mathbb{P}^1$ (French), Asterisque **86** (1981), 97-108.

[CH88] M. Cornalba, J. Harris, *Divisor classes associated to families of stable varieties with application to the moduli space of curves*, Ann. Sci. Ec. Norm. Sup., **21** (1988), 455-475.

[Fal83] G. Faltings, *Arakelov's theorem for abelian varieties*, Invent. Math. 73 (1983), no. 3, 337-347.

[Liu96] K. F. Liu, *Geometric height inequalities*, Math. Research Letters, **3** (1996), 637-702.

[Tan94] S.-L. Tan, *On the base changes of penciles of curves*, I, Manus. Math., **84** (1994), 225–244.

[Tan95] S.-L. Tan, *The minimal number of singular fibers of a semistable curve over* $\mathbb{P}^1$, J. Alg. Geom., **4** (1995), 591-596.

[VZ01] E. Viehweg, K. Zuo, *On the isotriviality of families of projective manifolds over curves*, J. Alg. Geom., **10** (2001) 781-799.

[VZ06] E. Viehweg, K. Zuo, *Numerical bounds for semi-stable families of curves or of certain higher dimensional manifolds*, J. Alg. Geom., **15** (2006) 771-791.

[Voj88] P. Vojta, *Diophantine inequalities and Arakelov theory.* In: Lang, S., Introduction to Arakelov theory. Springer-Verlag, Berlin-Heidelberg -New York, 1988, pp. 155-178.

[Xia85] G. Xiao, *Surfaces fibrées en courbes degenre deux*, Lect. Notes in Math., **1137** (1985) Springer-Verlag.

[Xia87] G. Xiao, *Fibred algebraic surfaces with low slope*, Math. Ann., **276** (1987), 449-466.

[Yu00] F. Yu, *Note on families of semistable curves over* $\mathbb{P}^1$ *with* 4 *singular fibers whose Jacobian are non-compact*, Sci. in China, **53** (2010), no.7, 1711-1714.

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, DONGCHUAN RD 500, SHANGHAI 200241, P. R. OF CHINA
E-mail address: `jlu@math.ecnu.edu.cn`

DEPARTMENT OF MATHEMATICS, EAST CHINA NORMAL UNIVERSITY, DONGCHUAN RD 500, SHANGHAI 200241, P. R. OF CHINA
E-mail address: `sltan@math.ecnu.edu.cn`

DEPARTMENT OF MATHEMATICS, XIAMEN UNIVERSITY, XIAMEN 361005, P. R. OF CHINA
E-mail address: `vvyufei@gmail.com`

UNIVERSITÄT MAINZ, FACHBEREICH 08-PHYSIK MATHEMATIK UND INFORMATIK, 55099 MAINZ, GERMANY
E-mail address: `zuok@uni-mainz.de`